

CLINICAL PRACTICE GUIDELINE FOR: PALLIATIVE SEDATION

EFFECTIVE DATE: 2009; REVIEW DATE: 2012; RELATED POLICIES: Consent for Treatment Policy #111-1; RELATED CPG'S OR CLINICAL PATHWAYS:

Clinical Practice Guidelines are to be used in conjunction with clinical judgment to support clinician and patient decisions for best care and treatment.

1.0 PURPOSE

To establish ethically acceptable criteria and guidelines for the use of palliative sedation as a form of treatment for intractable symptoms (most common symptoms include agitated delirium, dyspnea, pain, bleeding) (sedation for psycho-existential distress is controversial (*Palliative Sedation guidelines – AHS – Capital zone*) associated with acute or chronic morbidity in the final stages of end of life care.

To support and optimize quality decision-making processes for patients, families and clinicians associated with the ethical and responsible application of palliative sedation therapy

2.0 DEFINITIONS

2.1 Palliative sedation is: The process of inducing and maintaining deep sleep, in the final hours to days of life, for the relief of severe suffering caused by one or more intractable symptoms when all appropriate alternative interventions have failed to bring adequate symptom relief.

Palliative Sedation is NOT, and should be distinguished from:

2.1.1 Temporary use of complete sedation with the intention of reversal within a limited time

2.1.2 Use of benzodiazepines, neuroleptics, and other drugs for the management of symptoms. (e.g., delirium, dyspnea, etc.), where their use is to provide partial sedation (Chater, 1996; Fine, 2001; Walton & Weistein, 2002)

2.2 Refractory or intractable symptom: A symptom is considered refractory if it cannot be adequately controlled despite aggressive therapy that does not compromise consciousness. A refractory symptom must be distinguished from the “difficult to manage symptom” in that a “difficult symptom” could respond, within a tolerable time frame, to aggressive interventions that yield adequate relief and preserve consciousness, without excessive adverse results.

Decision points to determine a symptom as refractory - Further invasive or noninvasive interventions are either:

1. Incapable of providing adequate relief

2. Associated with excessive and intolerable acute or chronic morbidity, or
3. Unlikely to provide relief within a tolerable time frame (Cherny and Portenoy, 1994)

3.0 RATIONALE

- 3.1 The use of palliative sedation has been confused with euthanasia by various members of the health care team and public, leading to a need to clarify the indications for use. This guideline does not include any support for the practice of euthanasia.
- 3.2 The use of palliative sedation likely **does not** hasten death (Maltoni, 2009; Cherny, 2009), but further studies are needed.
- 3.3 The guidelines and decision-making algorithm for palliative sedation will enable clinicians to recognize patients who may be candidates for palliative sedation.
- 3.4 An approved guideline outlining the practice of palliative sedation may provide the staff with organizational support and guidance to improve comfort with initiating and administering palliative sedation.
- 3.5 The palliative sedation guideline provides a common base of understanding for what palliative sedation is and is not. Further, it allows patients, families, clinicians and care providers to make appropriate and informed choices.

4.0 CRITERIA

The basic criteria for considering the use of palliative sedation include all of the following;

- A terminal disease exists
- The patient/client suffers from (a) refractory symptom(s)
- In all but the most unusual circumstances, death must be imminent (within days)
- C2 goals of care designation (“*Goals of Care and interventions are for physical, psychological and spiritual preparation for imminent death (usually within hours or days). Maximal efforts directed at compassionate symptom control. Transfer is usually not undertaken*”) must be in effect.
- Informed consent must be obtained from the patient or substitute decision maker (as outlined in an Advanced Directive or a relative from a ranked list). If the patient is incapable or a substitute decision maker is not available, 2 physicians documented consent will also suffice (Alberta Adult Guardianship and Trusteeship Act).
- Palliative care physician input must be obtained

5.0 PROCESS

- 5.1 In considering the use of palliative sedation, the attending physician shall ensure the patient is assessed by an interdisciplinary palliative care team that includes, at minimum, a palliative care physician. There may be circumstances where a telephone or video-consultation will be considered appropriate as an interim measure or where geographical circumstances preclude a face to face consultation. In rural areas where a face to face consultation is not immediately possible, the patient should be discussed with the Rural Palliative Physician or with the Palliative Physician on-call if after hours.
- 5.2 The attending physician and the palliative care physician/team shall consult directly with the patient and family and, as appropriate, with the other care providers regarding the option of palliative sedation.

5.3 The physician shall ensure that the discussion by which the appropriate consent was obtained is documented on the health record. This documentation should include mention of all criteria met.

5.4 Once consent is obtained for palliative sedation, the palliative care physician/team will arrange for palliative sedation and appropriate monitoring of the patient.

6.0 SCOPE AND LIMITATIONS

6.1 The expertise required to assess for the suitability of palliative sedation is a specialized skill set requiring advanced education and experience in hospice palliative care medicine.

6.2 This guideline is intended to apply to patients as identified by the palliative care physician as suitable for palliative sedation.

6.3 In all but the most unusual circumstances, death must appear imminent (within days).

7.0 PALLIATIVE SEDATION ORDERING AND MONITORING REQUIREMENTS

7.1 Whenever possible, palliative sedation is to be ordered by a Palliative Care Physician. In situations where a telephone or video-consultation has determined that palliative sedation is appropriate, the attending physician may order palliative sedation as discussed with the Palliative Care Physician. Please make note of section 5.1.

7.2 Monitoring of patient response to palliative sedation and any required adjustments to palliative sedation delivery rate and amount (within the range detailed in the palliative physician's orders) will be by a registered nurse, optimally with experience in palliative sedation delivery. If the patient is at home, then the family can be trained by the registered nurse to monitor and adjust the medication administration, under close supervision (both face-to-face and phone) of that registered nurse.

7.3 It is recommended that a patient response monitoring tool (eg Riker scale (see Appendix)) be used to ensure that all care providers are aware of the degree of patient sedation that was agreed on.

8.0 GOAL OF MONITORING PALLIATIVE SEDATION (as per definition):

Inducing and maintaining deep sleep, in the final hours to days of life, for the relief of severe suffering.

8.1 Key observations and assessment indicating this goal is not being achieved is a trigger to clinical decision for adjusting palliative sedation.

9.0 DOCUMENTATION:

Documentation should include:

- “This is palliative sedation”
- The discussion with the patient and family by which all criteria were met, and the
- appropriate consent was obtained
- Name and dosage of medication administered
- Changes to dose of medication

- Patient's level of sedation (e.g. Riker scale goal)
- Nature of information and support offered to patient and/or family

SITE SPECIFIC REQUIREMENTS: Palliative sedation can be undertaken in any location (home, hospital or care centre) provided that the above criteria can be satisfied.

Background Information Alberta Health Services - Calgary Zone Clinical Practice Guideline for Palliative Sedation

Note about the evidence used in the guideline:

This document has been developed by incorporating the existing literature on the use of palliative sedation with the experiences of multidisciplinary pain and symptom management experts to produce a consensus based guideline.

REFERENCES

Adult Guardianship and Trusteeship Act, Legislative Assembly of Alberta, 2009.

Personal Directives Act, Legislative Assembly of Alberta, 2008.

Chater, S. (1996). Terminal Sedation Workshop. 11th International Congress on Care of the Terminally Ill. Montreal, Canada.

Chater, S., Viola, R., Paterson, J., Jarvis, V. (1998). Sedation for Intractable Distress in the Dying—A Survey of Experts. Palliative Medicine, 12, 255-269.

Cherny, N.I., Portenoy, R.K. (1994). Sedation in the Management of Refractory Symptoms: Guidelines for Evaluation and Treatment. Journal of Palliative Care, 10(2), 31-38.

Cherny, N. (2009). The use of sedation to relieve cancer patients' suffering at the end of life: addressing critical issues. Annals of Oncology, 20: 1153-1155.

Fine, P. (2001). Total sedation in end of life care: Clinical considerations. Journal of Hospice and Palliative Nursing. 3(3), 81-87.

Greene, W.R., Davis, W.H. (1991). Titrated Intravenous Barbiturates in the Control of Symptomatic Patients with Terminal Cancer. Southern Medical Journal, 84(3), 32-37.

Maltoni, M., Pittureri, C., Scarpi, E., Piccinini, L., Martin, F., Turci, P., Montanari, L., Nanni, O., Amadori, D. (2009). Palliative sedation therapy does not hasten death: results from a prospective multicenter study. Annals of Oncology, 20: 1163-1169.

Palliative Sedation guidelines, Alberta Health Services – Capital Zone, 2005 (<http://www.palliative.org/PC/ClinicalInfo/Clinical%20Practice%20Guidelines/PDF%20files/3A6%20Palliative%20Sedation%20and%20Addendum.pdf>)

Roy, R.J. (1990). Euthanasia – Taking a Stand. Journal of Palliative Care 6(1), 3-5.

Walton, O. & Weistein, S. (2002). Sedation and comfort at the end of life. Current Pain and Headache Reports, 6(3), 197-201.

AUTHORS

This Clinical Practice Guideline for Palliative Sedation was developed by the Clinical Practice Committee of the Palliative and End of Life Care, Alberta Health Services – Calgary Zone. For more information, please contact:

Medical Leader, Palliative and End of Life Care
Alberta Health Services – Calgary Zone
Telephone: (403) 944-2304
Fax: (403) 270-9652

COMMITMENT TO CPG MANAGEMENT

Ongoing management of the clinical practice guideline includes implementation, evaluation, and revisions for the purpose of maintaining a current evidence based document to guide the delivery of palliative sedation within the AHS – Calgary Zone. The responsibility for upkeep of this document rests within the authoring clinical team and should be clearly named here within this document as the current Medical Director, Palliative & Hospice Care Services. Guidelines should be reviewed at a minimum of every 3 years or as new evidence or other guidelines become available.

Palliative Sedation Guideline - Developed 2005; Revised 2009; Review Due 2012

Appendix

Riker Sedation – Agitation Scale

Score	Term	Descriptor
7	Dangerous Agitation	Pulling at IV/SC tubing, trying to remove catheters, climbing over bedrail, striking at staff, thrashing side-to-side.
6	Very agitated	Does not calm despite frequent, verbal reminding of limits, requires physical restraints.
5	Agitated	Anxious or mildly agitated, attempting to sit up, calms down to verbal instructions.
4	Calm and cooperative	Calm, awakens easily, follows commands.
3	Sedated	Difficult to arouse, awakens to verbal stimuli or gentle shaking but drifts off again, follows simple commands.
2	Very sedated	Arouses to physical stimuli but does not communicate or follow commands, may move spontaneously.
1	Unarousable	Minimal or no response to noxious stimuli, does not communicate or follow commands.